

A KADISON–SAKAI TYPE THEOREM

MADJID MIRZAVAZIRI AND MOHAMMAD SAL MOSLEHIAN

ABSTRACT. The celebrated Kadison–Sakai theorem states that every derivation on a von Neumann algebra is inner. In this paper, we prove this theorem for ultraweakly continuous $*$ - σ -derivations, where σ is an ultraweakly continuous surjective $*$ -linear mapping. We decompose a σ -derivation into a sum of an inner σ -derivation and a multiple of a homomorphism. The so-called $*(\sigma, \tau)$ -derivations are also discussed.

1. INTRODUCTION

One of important questions in the theory of derivations is that “When are all bounded derivations inner?” Forty years ago, R.V. Kadison [4] and S. Sakai [8] independently proved that every derivation of a von Neumann algebra $\mathfrak{M}$ into itself is inner. This was the starting point for the study of the so-called bounded cohomology groups. This nice result can be restated as saying that the first bounded cohomology group $H^1(\mathfrak{M}; \mathfrak{M})$ (i.e. the vector space of derivations modulo the inner derivations) vanishes. One may conjecture the vanishing of the higher order bounded cohomology groups $H^n(\mathfrak{M}; \mathfrak{M}), n \geq 2$. It is remarkable to note that if $\mathfrak{M}$ is a von Neumann algebra of type I, II_1 , III , or is of type II_1 and stable under tensoring with the hyperfinite II_1 factor, then $H^n(\mathfrak{M}; \mathfrak{M}) = 0$ and $H^n(\mathfrak{M}; \mathcal{B}(\mathfrak{H})) = 0$ for all $n \geq 1$. Not all type II_1 factors are stable under tensoring with the hyperfinite II_1 factor, and so this theorem leaves open the type II_1 case. It is still unknown whether $H^1(\mathfrak{M}, \mathcal{B}(\mathfrak{H}))$ is always zero; see [9, 10].

Suppose that $\mathfrak{A}$ and $\mathfrak{B}$ are two algebras, $\mathfrak{X}$ is a $\mathfrak{B}$ -bimodule and $\sigma : \mathfrak{A} \rightarrow \mathfrak{B}$ is a linear mapping, which is called the ground mapping. A linear mapping $d : \mathfrak{A} \rightarrow \mathfrak{X}$ is called a σ -derivation if $d(ab) = d(a)\sigma(b) + \sigma(a)d(b)$ for all $a, b \in \mathfrak{A}$. These maps have been extensively investigated in pure algebra. Recently, they have been treated in the Banach algebra theory (see [1, 3, 5, 6, 7, 12] and references therein).

2000 *Mathematics Subject Classification.* Primary 46L57; Secondary 46L05, 47B47.

Key words and phrases. Derivation, $*$ -homomorphism, $*$ - σ -derivation, inner σ -derivation, $*(\sigma, \tau)$ -derivation, Kadison–Sakai theorem, ultraweak (operator) topology.

There are some applications of σ -derivations to develop an approach to deformations of Lie algebras which have many applications in models of quantum phenomena and in analysis of complex systems; cf. [2]. A wide range of examples are as follows:

- (i) Every ordinary derivation of an algebra $\mathfrak{A}$ into an $\mathfrak{A}$ -bimodule $\mathfrak{X}$ is an $\iota_{\mathfrak{A}}$ -derivation (throughout the paper, $\iota_{\mathfrak{A}}$ denotes the identity map on the algebra $\mathfrak{A}$);
- (ii) Every endomorphism α on $\mathfrak{A}$ is a $\frac{\alpha}{2}$ -derivation;
- (iii) For a given homomorphism ρ on $\mathfrak{A}$ and a fixed arbitrary element X in an $\mathfrak{A}$ -bimodule $\mathfrak{X}$, the linear mapping $\delta(A) = X\rho(A) - \rho(A)X$ is a ρ -derivation of $\mathfrak{A}$ into $\mathfrak{X}$ which is said to be an inner ρ -derivation.
- (iv) Every point derivation $d : \mathfrak{A} \rightarrow \mathbb{C}$ at the character θ is a (θ, θ) -derivation.

In this paper, we investigate σ -derivations. We divide our main work into three sections. The first section is devoted to prove a Kadison–Sakai type theorem for ρ -derivations on von Neumann algebras when ρ is a homomorphism. In the next section, section we briefly discuss an extension of one of our result to the (σ, τ) -derivations. In the last section, we decompose a σ -derivation into a sum of an inner σ -derivation and a multiple of a homomorphism. The importance of our approach is that σ is a linear mapping in general, not necessarily a homomorphism.

A von Neumann algebra $\mathfrak{M}$ is an ultraweakly closed $*$ -subalgebra of $B(\mathfrak{H})$ containing the identity operator I , where $\mathfrak{H}$ is a Hilbert space. By the weak (operator) topology on $B(\mathfrak{H})$ we mean the topology generated by the semi-norms $T \mapsto |\langle T\xi, \eta \rangle|$ ($\xi, \eta \in \mathfrak{H}$). We also use the terminology ultraweak (operator) topology for the weak*-topology on $B(\mathfrak{H})$ considered as the dual space of the nuclear operators on $\mathfrak{H}$. When we speak of ultraweak continuity (weak continuity, resp.) of a mapping between von Neumann algebras $\mathfrak{M}$ and $\mathfrak{N}$ we mean that we equipped both $\mathfrak{M}$ and $\mathfrak{N}$ with the ultraweak topology (the weak topology, resp.). We refer the reader to [11] for undefined phrases and notations.

2. TOWARDS A KADISON–SAKAI TYPE THEOREM FOR σ -DERIVATIONS

We start our work with the following lemma.

Lemma 2.1 (S. Sakai). *Let $\rho : \mathfrak{M} \rightarrow \mathfrak{N}$ be an ultraweakly continuous $*$ -epimorphism and $\delta : \mathfrak{M} \rightarrow \mathfrak{N}$ be a $*$ - ρ -derivation between von Neumann algebras. Then there is a central projection $P \in \mathfrak{M}$ and an $*$ -isomorphism $\tilde{\rho} : \mathfrak{M}P \rightarrow \mathfrak{N}$ such that $\delta : \mathfrak{M}P \rightarrow \mathfrak{N}$ is a $*$ - $\tilde{\rho}$ -derivation. Moreover, $\delta(A) = 0$ for each $A \in \mathfrak{M}(I - P)$. Also, $\tilde{\rho}^{-1} \circ \delta$ is an ordinary derivation on $\mathfrak{M}P$.*

Proof. Since ρ is an ultraweakly continuous $*$ -homomorphism, its kernel is an ultraweakly closed ideal of $\mathfrak{M}$. Hence there is a central projection $Q \in \mathfrak{M}$ such that $\ker \rho = \mathfrak{M}Q$. Set $P = I - Q$. Then $\tilde{\rho} = \rho|_{\mathfrak{M}P} : \mathfrak{M}P \rightarrow \mathfrak{N}$ is a $*$ -isomorphism. We have

$$\delta(ABP) = \delta(APBP) = \delta(AP)\rho(AP) + \rho(AP)\delta(BP).$$

Hence δ is a $*$ - $\tilde{\rho}$ -derivation on $\mathfrak{M}P$. Moreover, if $A = B(I - P) \in \mathfrak{M}(I - P)$ is a projection then

$$\delta(A) = \delta((BQ)^2) = \delta(BQ)\rho(BQ) + \rho(BQ)\delta(BQ) = 0,$$

since $BQ \in \mathfrak{M}Q = \ker \rho$. The space $\mathfrak{M}(I - P)$ is a von Neumann algebra, because it is $\ker \rho = \rho^{-1}(\{0\})$ and ρ is ultraweakly continuous. Thus $\mathfrak{M}(I - P)$ is generated by its projection and so $\delta(A) = 0$ for each $A \in \mathfrak{M}(I - P)$. The last assertion is now obvious. $\square$

Theorem 2.2. *If $\rho : \mathfrak{M} \rightarrow \mathfrak{N}$ is an ultraweakly continuous $*$ -epimorphism and $\delta : \mathfrak{M} \rightarrow \mathfrak{N}$ be an ultraweakly continuous $*$ - ρ -derivation between von Neumann algebras, then there is an element $U \in \mathfrak{M}$ with $\|U\| \leq \|\delta\|$ such that $\delta(A) = U\rho(A) - \rho(A)U$ for each $A \in \mathfrak{M}$. In other words, δ is an inner ρ -derivation.*

Proof. The mapping $\tilde{\rho}^{-1} \circ \delta$ is an ordinary derivation on $\mathfrak{M}P$, where P is as in Lemma 2.1. Thus, by [8, Theorem 2.5.3], there is a $V \in \mathfrak{M}P$ with $\|V\| \leq \|\tilde{\rho}^{-1} \circ \delta\| \leq \|\delta\|$ such that $(\tilde{\rho}^{-1} \circ \delta)(A) = VA - AV$ for all $A \in \mathfrak{M}P$. Thus $\delta(A) = \tilde{\rho}(V)\tilde{\rho}(A) - \tilde{\rho}(A)\tilde{\rho}(V)$, for all $A \in \mathfrak{M}P$. Putting $U = \tilde{\rho}(V)$ we have $\delta(A) = U\rho(A) - \rho(A)U$ for all $A \in \mathfrak{M}P$. The later equality is also valid for $A \in \mathfrak{M}(I - P)$, since both sides are 0 for these elements. Finally, $\|U\| = \|\rho(V)\| \leq \|V\| \leq \|\delta\|$. $\square$

The following example can illustrate the case that the ground mapping is not a homomorphism.

Example 2.3. Let $\mathfrak{H}$ be an infinite dimensional Hilbert space with an orthonormal basis $\{e_n\}_{n \in \mathbb{N}}$, and $\theta : \mathcal{B}(\mathfrak{H}) \rightarrow \mathbb{C}$ be a non-zero character. Define $\rho : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ by

$$\langle \rho(A)e_n, e_m \rangle = \begin{cases} \theta(A) & \text{if } n = m = 1 \\ 0 & \text{otherwise} \end{cases}$$

and $\delta : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ by

$$\langle \delta(A)e_n, e_m \rangle = \begin{cases} \theta(A) & \text{if } (n, m) = (1, 2) \text{ or } (n, m) = (2, 1) \\ 0 & \text{otherwise} \end{cases}$$

A simple verification by using the matrix form of $\rho(A)$ and $\delta(A)$ shows that ρ is a $*$ -homomorphism which is not surjective and δ is a $*$ - ρ -derivation. Clearly, $\delta(I)$ does not commute with $\rho(I)$.

Let there be a non-zero $*$ - θ -derivation $\zeta : \mathcal{B}(\mathfrak{H}) \rightarrow \mathbb{C}$. Then $\delta : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ defined by

$$\langle \delta(A)e_n, e_m \rangle = \begin{cases} \zeta(A) & \text{if } n = m = 1 \\ \theta(A) & \text{if } (n, m) = (1, 2) \text{ or } (n, m) = (2, 1) \\ 0 & \text{otherwise} \end{cases}$$

is a $*$ - ρ -derivation which is not ρ -inner, is not a $*$ -homomorphism and does not decompose to a sum of a ρ -inner derivation and a $*$ -homomorphism.

3. (σ, τ) -DERIVATIONS

Assume that $\mathfrak{A}, \mathfrak{B}$ are $*$ -algebras and $\sigma, \tau : \mathfrak{A} \rightarrow \mathfrak{B}$ are $*$ -linear mappings. By a (σ, τ) -derivation we mean a linear mapping $d : \mathfrak{A} \rightarrow \mathfrak{B}$ preserving $*$ such that $d(ab) = d(a)\sigma(b) + \tau(a)d(b)$ for all $a, b \in \mathfrak{A}$. Obviously, a $*$ - σ -derivation is a (σ, σ) -derivation.

Proposition 3.1. *Let $\mathfrak{A}, \mathfrak{B}$ be $*$ -algebras and $\sigma, \tau : \mathfrak{A} \rightarrow \mathfrak{B}$ be $*$ -linear mappings. Then any (σ, τ) -derivation $d : \mathfrak{A} \rightarrow \mathfrak{B}$ is a $*$ - $\frac{\sigma+\tau}{2}$ -derivation.*

Proof. First we show that each (σ, τ) -derivation is a (τ, σ) -derivation. We have $d(ab) = d(b^*a^*)^* = (d(b)^*\sigma(a)^* + \tau(b)^*d(a)^*)^* = d(a)\tau(b) + \sigma(a)\tau(b)$.

Now we conclude that $d(ab) = \frac{1}{2}d(ab) + \frac{1}{2}d(ab) = \frac{1}{2}(d(a)\sigma(b) + \tau(a)d(b)) + \frac{1}{2}(d(a)\tau(b) + \sigma(a)d(b)) = d(a)\frac{\sigma+\tau}{2}(b) + \frac{\sigma+\tau}{2}(a)d(b)$. $\square$

The previous theorem enables us to focus on $*$ - σ -derivations while we deal with $*$ -algebras. In particular we obtain the following generalization of Theorem 2.2.

Theorem 3.2. *Let $\mathfrak{M}, \mathfrak{N}$ be von Neumann algebras, let $\rho_1, \rho_2 : \mathfrak{M} \rightarrow \mathfrak{N}$ be ultraweakly continuous and norm continuous $*$ -homomorphisms such that $\rho_1 + \rho_2$ be surjective, and let $\delta : \mathfrak{M} \rightarrow \mathfrak{N}$ be an ultraweakly continuous (ρ_1, ρ_2) -derivation. Then there is an element U in $\mathfrak{M}$ with $\|U\| \leq \frac{\|\rho_1\| + \|\rho_2\|}{2} \|\delta\|$ such that $\delta(A) = U \frac{\rho_1(A) + \rho_2(A)}{2} - \frac{\rho_1(A) + \rho_2(A)}{2} U$ for each $A \in \mathfrak{M}$.*

4. A KADISON-SAKAI TYPE THEOREM FOR σ -DERIVATIONS

In this section, we aim to remove the condition of being a homomorphism from the ground mapping and provide our main result on the generalization of the Kadison-Sakai

theorem. Indeed, we would like to decompose d as a direct sum of an inner $*$ - σ -derivation and a $*$ -homomorphism.

Recall that if we have a Hilbert space direct sum $\mathfrak{H} = \mathfrak{K} \oplus \mathfrak{L}$, P is the projection corresponding to $\mathfrak{K}$ and $T \in \mathcal{B}(\mathfrak{H})$, then the compression $T_{\mathfrak{K}}$ to $\mathfrak{K}$ is the operator $T_{\mathfrak{K}} \in \mathcal{B}(\mathfrak{K})$ defined by $T_{\mathfrak{K}}(k) = PT(k)$ ($k \in \mathfrak{K}$). Obviously, $PTP \leftrightarrow T_{\mathfrak{K}}$ is an isometric $*$ -isomorphism between von Neumann algebras $P\mathcal{B}(\mathfrak{H})P$ and $\mathcal{B}(\mathfrak{K})$. If $\sigma : \mathfrak{M} \rightarrow \mathfrak{M}$ is a linear mapping such that $\sigma(A)P = P\sigma(A)$ for all $A \in \mathfrak{M}$, then $\sigma_{\mathfrak{K}} : \mathfrak{M} \rightarrow P\mathfrak{M}P$, $\sigma_{\mathfrak{L}} : \mathfrak{M} \rightarrow (I - P)\mathfrak{M}(I - P)$ can be defined by $\sigma_{\mathfrak{K}}(A) = P\sigma(A)P$, $\sigma_{\mathfrak{L}}(A) = (I - P)\sigma(A)(I - P)$. Furthermore, $\sigma = \sigma_{\mathfrak{K}} \oplus \sigma_{\mathfrak{L}}$. Note that $P\mathfrak{M}P$ ($(I - P)\mathfrak{M}(I - P)$, resp.) is a von Neumann subalgebra of $B(\mathfrak{K})$ ($B(\mathfrak{L})$, resp.).

Example 4.1. Let $\mathfrak{H}$ be the separable Hilbert space with orthonormal basis $\{e_n\}$. Define $\sigma : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ by

$$\langle \sigma(A)e_n, e_m \rangle = \begin{cases} \exp(\frac{i}{s} - \frac{i}{r}) \langle Ae_s, e_r \rangle & \text{if } n = 2s - 1, m = 2r - 1 \\ \frac{1}{2} \exp(\frac{i}{s} - \frac{i}{r}) \langle Ae_s, e_r \rangle & \text{if } n = 2s, m = 2r \\ 0 & \text{otherwise} \end{cases}$$

and $d : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{H})$ by

$$\langle d(A)e_n, e_m \rangle = \begin{cases} (\frac{1}{s} - \frac{1}{r}) \exp(\frac{i}{s} - \frac{i}{r}) \langle Ae_s, e_r \rangle & \text{if } n = 2s - 1, m = 2r - 1 \\ \exp(\frac{i}{s} - \frac{i}{r}) \langle Ae_s, e_r \rangle & \text{if } n = 2s, m = 2r \\ 0 & \text{otherwise} \end{cases}$$

Then d is a σ -derivation. To show this let E_{rs} be the operator defined on $\mathfrak{H}$ by

$$\langle E_{rs}e_n, e_m \rangle = \delta_{rm} \delta_{sn},$$

where δ is the δ -Kronecker function. Then

$$\langle d(E_{rs})e_n, e_m \rangle = \begin{cases} (\frac{1}{s} - \frac{1}{r}) \exp(\frac{i}{s} - \frac{i}{r}) & \text{if } n = 2s - 1, m = 2r - 1 \\ \exp(\frac{i}{s} - \frac{i}{r}) & \text{if } n = 2s, m = 2r \\ 0 & \text{otherwise} \end{cases}$$

and so

$$\langle d(E_{tu}E_{rs})e_n, e_m \rangle = \delta_{ur} \begin{cases} (\frac{1}{s} - \frac{1}{t}) \exp(\frac{i}{s} - \frac{i}{t}) & \text{if } n = 2s - 1, m = 2t - 1 \\ \exp(\frac{i}{s} - \frac{i}{t}) & \text{if } n = 2s, m = 2t \\ 0 & \text{otherwise} \end{cases}$$

On the other hand,

$$\langle \sigma(E_{rs})e_n, e_m \rangle = \begin{cases} \exp(\frac{i}{s} - \frac{i}{r}) & \text{if } n = 2s - 1, m = 2r - 1 \\ \frac{1}{2} \exp(\frac{i}{s} - \frac{i}{r}) & \text{if } n = 2s, m = 2r \\ 0 & \text{otherwise} \end{cases}$$

and so

$$\langle d(E_{tu})\sigma(E_{rs})e_n, e_m \rangle = \delta_{ur} \begin{cases} (\frac{1}{u} - \frac{1}{t}) \exp(\frac{i}{s} - \frac{i}{t}) & \text{if } n = 2s - 1, m = 2t - 1 \\ \frac{1}{2} \exp(\frac{i}{s} - \frac{i}{t}) & \text{if } n = 2s, m = 2t \\ 0 & \text{otherwise} \end{cases}$$

and

$$\langle \sigma(E_{tu})d(E_{rs})e_n, e_m \rangle = \delta_{ur} \begin{cases} (\frac{1}{s} - \frac{1}{u}) \exp(\frac{i}{s} - \frac{i}{t}) & \text{if } n = 2s - 1, m = 2t - 1 \\ \frac{1}{2} \exp(\frac{i}{s} - \frac{i}{t}) & \text{if } n = 2s, m = 2t \\ 0 & \text{otherwise} \end{cases}$$

This shows that

$$d(E_{tu}E_{rs}) = d(E_{tu})\sigma(E_{rs}) + \sigma(E_{tu})d(E_{rs})$$

and so d is a σ -derivation.

Now let $\mathfrak{K}$ be the Hilbert space with orthonormal basis $\{e_{2n-1}\}$ and $\mathfrak{L}$ be the Hilbert space with orthonormal basis $\{e_{2n}\}$. Then $\mathfrak{H} = \mathfrak{K} \oplus \mathfrak{L}$. If $\sigma_{\mathfrak{K}} : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{K})$ and $\sigma_{\mathfrak{L}} : \mathcal{B}(\mathfrak{H}) \rightarrow \mathcal{B}(\mathfrak{L})$ are defined as above, then $\sigma = \sigma_{\mathfrak{K}} \oplus \sigma_{\mathfrak{L}}$. By the same fashion we can write $d = d_{\mathfrak{K}} \oplus d_{\mathfrak{L}}$. One can easily verify that $d_{\mathfrak{K}}$ is a $\sigma_{\mathfrak{K}}$ -derivation, $d_{\mathfrak{L}}$ is a $\sigma_{\mathfrak{L}}$ -derivation and $\sigma_{\mathfrak{K}}$ is a $*$ -homomorphism on unitary group, $\sigma_{\mathfrak{L}}$ is $\frac{1}{2}$ times a $*$ -homomorphism, and $d_{\mathfrak{K}}(A) = U\sigma_{\mathfrak{K}}(A) - \sigma_{\mathfrak{K}}(A)U$, where U is the operator defined by $\langle Ue_{2n-1}, e_{2m-1} \rangle = \frac{1}{n}\delta_{nm}$. Moreover $d_{\mathfrak{L}} = 2\sigma_{\mathfrak{L}}$. Note also that we have $\sigma_{\mathfrak{K}}(I) = I_{\mathfrak{K}}$, $\sigma_{\mathfrak{L}}(I) = \frac{1}{2}I_{\mathfrak{L}}$, $d_{\mathfrak{K}}(I) = 0$ and $d_{\mathfrak{L}}(I) = I_{\mathfrak{L}}$.

We will show that the situation described in the above example is true in general.

Proposition 4.2. *Suppose that $\sigma : \mathfrak{M} \rightarrow \mathfrak{M}$ is an ultraweakly continuous surjective $*$ -linear mapping, $d : \mathfrak{M} \rightarrow \mathfrak{M}$ is an ultraweakly continuous $*$ - σ -derivation such that $d(I)$ is a central element of $\mathfrak{M}$, $\mathfrak{L}_0 = \bigcup_{B,C \in \mathfrak{M}} (\sigma(BC) - \sigma(B)\sigma(C))(\mathfrak{H})$, $\mathfrak{L}$ is the closed linear span of $\mathfrak{L}_0$ and $\mathfrak{K} = \mathfrak{L}^\perp$. Then*

- (i) $\mathfrak{K} = \bigcap_{B,C \in \mathfrak{M}} \ker(\sigma(BC) - \sigma(B)\sigma(C))$;
- (ii) *If $P = P_{\mathfrak{K}}$ is the projection corresponding to $\mathfrak{K}$, then $\sigma(A)P = P\sigma(A)$ and $d(A)P = Pd(A)$ for all $A \in \mathfrak{M}$;*

(iii) If we define $\delta : \mathfrak{M} \rightarrow P\mathfrak{M}P$ by $\delta(A) = Pd(A)P$, $\rho : \mathfrak{M} \rightarrow P\mathfrak{M}P$ by $\rho(A) = \sigma(A)P$, $\alpha : \mathfrak{M} \rightarrow (I - P)\mathfrak{M}(I - P)$ by $\alpha(A) = (I - P)d(A)(I - P)$ and $\tau : \mathfrak{M} \rightarrow (I - P)\mathfrak{M}(I - P)$ by $\tau(A) = (I - P)\sigma(A)(I - P)$, then δ is an ultraweakly continuous $*$ - ρ -derivation, α is an ultraweakly continuous $*$ - τ -derivation, $d = \delta \oplus \alpha$ and $\sigma = \rho \oplus \tau$. Moreover ρ is an ultraweakly continuous $*$ -epimorphism;

(iv) $\mathfrak{K} = \ker \delta(I)$ and $\mathfrak{L} = \overline{\alpha(I)(\mathfrak{L})}$;

(v) δ is an inner ρ -derivation.

(vi) $\tau(I) = \frac{1}{2}I_{\mathfrak{L}}$

Proof. (i) For each $B, C \in \mathfrak{M}$, $h \in \mathfrak{H}$ and $k \in \mathfrak{K}$ we have

$$\begin{aligned} 0 &= \langle (\sigma(BC) - \sigma(B)\sigma(C))(h), k \rangle \\ &= \langle h, (\sigma(BC) - \sigma(B)\sigma(C))^*(k) \rangle \\ &= \langle h, (\sigma(C^*B^*) - \sigma(C^*)\sigma(B^*))(k) \rangle. \end{aligned}$$

Since $\mathfrak{M}$ is a $*$ -subalgebra of $\mathcal{B}(\mathfrak{H})$, we infer that $(\sigma(BC) - \sigma(B)\sigma(C))(k) = 0$ for each $B, C \in \mathfrak{M}$ and $k \in \mathfrak{K}$. This shows that $\mathfrak{K} = \bigcap_{B, C \in \mathfrak{M}} \ker(\sigma(BC) - \sigma(B)\sigma(C))$.

(ii) Let $P = P_{\mathfrak{K}}$ be the projection corresponding to $\mathfrak{K}$. For each $B, C, A \in \mathfrak{M}$ and $k \in \mathfrak{K}$ we have

$$\begin{aligned} (\sigma(BC) - \sigma(B)\sigma(C))\sigma(A)(k) &= (\sigma(BC)\sigma(A) - \sigma(B)\sigma(C)\sigma(A))(k) \\ &= (\sigma(BCA) - \sigma(BCA))(k) \\ &= 0. \end{aligned}$$

This shows that $\sigma(A)(\mathfrak{K}) \subseteq \mathfrak{K}$ for each $A \in \mathfrak{M}$. Since σ is a $*$ -linear mapping, we have $\sigma(A)P = P\sigma(A)$ for all $A \in \mathfrak{M}$.

By using [6, Lemma 2.2] we get

$$0 = d(B)(\sigma(CA) - \sigma(C)\sigma(A))(k) = (\sigma(BC) - \sigma(B)\sigma(C))d(A)(k)$$

for all $k \in \mathfrak{K}$. This implies that $d(A)(k) \in \mathfrak{K}$ for all $k \in \mathfrak{K}$. Hence $d(A)(\mathfrak{K}) \subseteq \mathfrak{K}$ for all $A \in \mathfrak{M}$. Since d is a $*$ -linear mapping we conclude that $d(A)P = Pd(A)$ for all $A \in \mathfrak{M}$.

(iii) Firstly, ρ is a $*$ -homomorphism. In fact, if $B, C \in \mathfrak{M}$ then

$$\begin{aligned}
 \rho(BC)(k) &= \sigma(BC)P(k) \\
 &= \sigma(BC)(k) \\
 &= \sigma(B)\sigma(C)(k) \\
 &= \sigma(B)\sigma(C)P^2(k) \\
 &= \sigma(B)P\sigma(C)P(k) \\
 &= \rho(B)\rho(C)(k)
 \end{aligned}$$

for all $k \in \mathfrak{K}$, and

$$\rho(BC)(\ell) = \sigma(BC)P(\ell) = 0 = \sigma(B)P\sigma(C)P(\ell) = \rho(B)\rho(C)(\ell).$$

for all $\ell \in \mathfrak{L}$. Hence ρ is a homomorphism on $\mathfrak{M}$. Moreover, $\rho(A^*) = \sigma(A^*)P = \sigma(A)^*P = (P\sigma(A))^* = (\sigma(A)P)^* = \rho(A)^*$, ($A \in \mathfrak{M}$).

Secondly, δ is a ρ -derivation, since if $A, B \in \mathfrak{M}$ then

$$\begin{aligned}
 \delta(AB)(k) &= d(AB)P(k) \\
 &= Pd(AB)(k) \\
 &= Pd(A)\sigma(B)(k) + P\sigma(A)d(B)(k) \\
 &= P^2d(A)\sigma(B)P^2(k) + P^2\sigma(A)d(B)P^2(k) \\
 &= Pd(A)PP\sigma(B)P(k) + P\sigma(A)PPd(B)P(k) \\
 &= (\delta(A)\rho(B) + \rho(A)d(B))(k),
 \end{aligned}$$

for all $k \in \mathfrak{K}$, and

$$\begin{aligned}
 \delta(AB)(\ell) &= Pd(AB)P(\ell) \\
 &= 0 \\
 &= Pd(A)PP\sigma(B)P(\ell) + P\sigma(A)PPd(B)P(\ell) \\
 &= (\delta(A)\rho(B) + \rho(A)d(B))(\ell).
 \end{aligned}$$

for all $\ell \in \mathfrak{L}$.

Similarly one can show that α is a τ -derivation. It is obvious that $d = \delta \oplus \alpha$ and $\sigma = \rho \oplus \tau$.

(iv) We have

$$d(I) = d(I^2) = d(I)\sigma(I) + \sigma(I)d(I) = 2\sigma(I)d(I).$$

Since σ is surjective, $d(I)$ is of the form $\sigma(E)$ for some $E \in \mathfrak{M}$. Now for each $k \in \mathfrak{K}$ we have

$$d(I)(k) = \sigma(E)(k) = \sigma(EI)(k) = \sigma(E)\sigma(I)(k) = d(I)\sigma(I)(k)$$

and so $d(I)(k) = d(I)(2\sigma(I) - I)(k) = 0$ for each $k \in \mathfrak{K}$. Hence $\delta(I)(k) = d(I)P(k) = 0$ for each $k \in \mathfrak{K}$. Thus $k \subseteq \ker \delta(I)$. On the other hand the compression operator $\delta(I)$ belongs to $\mathcal{B}(\mathfrak{K})$. Hence $\ker \delta(I) \subseteq \mathfrak{K}$. Thus $\mathfrak{K} = \ker \delta(I)$. Similarly one can show that $\mathfrak{L} = \overline{\alpha(I)(\mathfrak{L})}$.

(v) By Theorem 2.2 there is an element $U \in \mathfrak{M}$ such that $\delta(A) = U\rho(A) - \rho(A)U$ for all $A \in \mathfrak{M}$. Hence δ is an inner ρ -derivation.

(vi) Since $d(I)$ is a central element of $\mathfrak{M}$, one easily deduce that $\alpha(I) = d(I)(I - P)$ is in the center of $\mathfrak{M}$. Hence

$$\alpha(I) = \alpha(I^2) = \alpha(I)\tau(I) + \tau(I)\alpha(I) = 2\tau(I)\alpha(I).$$

Thus $(2\tau(I) - I)\alpha(I) = 0$. It follows from $\overline{\alpha(I)(\mathfrak{L})} = \mathfrak{L}$ that $2\tau(I) = I_{\mathfrak{L}}$. □

We can now establish a version of the Kadison–Sakai theorem as our main theorem.

Theorem 4.3. *Suppose that $\sigma : \mathfrak{M} \rightarrow \mathfrak{M}$ is an ultraweakly continuous surjective $*$ -linear mapping and $d : \mathfrak{M} \rightarrow \mathfrak{M}$ is an ultraweakly continuous $*$ - σ -derivation such that $d(I)$ is a central element of $\mathfrak{M}$. Then $\mathfrak{H}$ can be decomposed into $\mathfrak{K} \oplus \mathfrak{L}$ and d can be factored as the form*

$$\delta \oplus 2Z\tau,$$

where $\delta : \mathfrak{M} \rightarrow \mathfrak{M}$ is an inner $*$ - $\sigma_{\mathfrak{K}}$ -derivation, Z is a central element, and $2\tau = 2\sigma_{\mathfrak{L}}$ is a $*$ -homomorphism.

Proof. It remains to show the result concerning $\mathfrak{L}$. For $\ell \in \mathfrak{L}$ we have

$$\alpha(A)(\ell) = \alpha(I)\tau(A)(\ell) + \tau(I)\alpha(A)(\ell) = \alpha(I)\tau(A)(\ell) + \frac{1}{2}\alpha(A)(\ell).$$

Thus

$$\alpha(A)(\ell) = 2\alpha(I)\tau(A)(\ell).$$

Putting $Z = \alpha(I)$ we get the result. □

REFERENCES

1. M. Brešar and A.R. Villena, *The noncommutative Singer-Werner conjecture and ϕ -derivations*, J. London Math. Soc. **(2) 66** (2002), no. 3, 710–720.
2. J. Hartwig, D. Larsson and S. D. Silvestrov, *Deformations of Lie algebras using σ -derivations*, Preprints in Math. Sci. 2003:32, LUTFMA-5036-2003 Centre for Math. Sci., Dept of Math., Lund Inst. of Tech., Lund Univ., 2003.
3. S. Hejazian, A.R. Janfada, M. Mirzavaziri and M.S. Moslehian, *Achievement of continuity of (φ, ψ) -derivations without linearity*, Bull. Belg. Math. Soc.-Simon Stevn., **14** (2007), no. 4, 641–652.
4. R.V. Kadison, *Derivations of operator algebras*, Ann. Math. **83** (1966), 280–293.
5. M. Mirzavaziri and M.S. Moslehian, *σ -derivations in Banach algebras*, Bull. Iranian Math. Soc. **32** (2006), no. 1, 65–78.
6. M. Mirzavaziri and M.S. Moslehian, *Automatic continuity of σ -derivations in C^* -algebras*, Proc. Amer. Math. Soc., **134** (2006), no. 11, 3319–3327.
7. M.S. Moslehian, *Approximate $(\sigma - \tau)$ -contractibility*, Nonlinear Funct. Anal. Appl., **11** (2006), no. 5, 805–813.
8. S. Sakai, *Derivations of W^* -algebras*, Ann. Math. **83** (1966), 273–279.
9. A.M. Sinclair and R.R. Smith, *Hochschild Cohomology of von Neumann Algebras*, Cambridge Univ. Press, Cambridge, 1995.
10. A.M. Sinclair and R.R. Smith, *A survey of Hochschild cohomology for von Neumann algebras*, Operator algebras, quantization, and noncommutative geometry, 383–400, Contemp. Math., 365, Amer. Math. Soc., Providence, RI, 2004.
11. M. Takesaki, *Theory of Operator Algebras. I*, Reprint of the first (1979) edition. Encyclopaedia of Mathematical Sciences. 124. Operator Algebras and Non-commutative Geometry, 5. Springer-Verlag, Berlin, 2002.
12. J.M. Zhan and Z.S. Tan, *T -local derivations of von Neumann algebras*, Northeast. Math. J. **20** (2004), no. 2, 145–152.

DEPARTMENT OF MATHEMATICS, FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASHHAD 91775, IRAN;

BANACH MATHEMATICAL RESEARCH GROUP (BMRG), MASHHAD, IRAN;

CENTRE OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIVERSITY OF MASHHAD, IRAN.

E-mail address: mirzavaziri@math.um.ac.ir and madjid@mirzavaziri.com

E-mail address: moslehian@ferdowsi.um.ac.ir and moslehian@ams.org